

Field induced stationary state for an accelerated tracer in a bath

Matthieu Barbier and Emmanuel Trizac

*Laboratoire de Physique Théorique et Modèles Statistiques,
UMR CNRS 8626, Université Paris-Sud, 91405 Orsay, France*

Our interest goes to the behavior of a tracer particle, accelerated by a constant and uniform external field, when the energy injected by the field is redistributed through collision to a bath of unaccelerated particles. A non equilibrium steady state is thereby reached. Solutions of a generalized Boltzmann-Lorentz equation are analyzed analytically, in a versatile framework that embeds the majority of tracer-bath interactions discussed in the literature. These results –mostly derived for a one dimensional system– are successfully confronted to those of three independent numerical simulation methods: a direct iterative solution, Gillespie algorithm, and the Direct Simulation Monte Carlo technique. We work out the diffusion properties as well as the velocity tails: large v , and either large $-v$, or v in the vicinity of its lower cutoff whenever the velocity distribution is bounded from below. Particular emphasis is put on the cold bath limit, with scatterers at rest, which plays a special role in our model.

I. INTRODUCTION

The propagation of a single test particle (or "tracer") with mass m_t accelerated by an external field g in a stationary bath of particles with mass m_b was first studied by Lorentz in 1905 [1] as a kinetic approach of conduction in a metal. In this model, an electric field accelerates non-interacting electrons, which diffuse among a random configuration of static obstacles - the atoms - acting as infinitely massive scatterers ($m_t/m_b \rightarrow 0$), with a simple hard sphere electron-atom collision rule. Lorentz assumed the velocity distribution of the electrons to obey a Maxwell-Boltzmann law with a small correction, and was able to give an estimate of the conductivity of metals from microscopic parameters. It was later shown that this perturbed Maxwell-Boltzmann distribution cannot be a stationary solution of the problem, although it displays the correct value for the mean velocity [2]: in the Lorentz model, the constant acceleration confers an ever-increasing amount of energy to the electron, which cannot be dissipated in the described collision process; Its velocity variance consequently diverges over time, despite the mean remaining finite. However, a stationary solution may exist if the tracer is allowed to transfer some energy to the surrounding medium, for instance if m_t/m_b is finite. The case $m_b = m_t$ was therefore subsequently considered [3], as a non-vanishing g extension of Résibois' "self-diffusion problem" [4], wherein one studies the diffusive properties of identical gas particles at equilibrium by following a single (unaccelerated) tracer. The bulk of ulterior developments on this modified Lorentz problem focused on asymptotic and relaxation properties [5–10]. Recently, the model was used as a paradigm for the study of microscopic entropy production and fluctuation relations in non equilibrium systems [11, 12], and to discuss the force-velocity relation [13] as measured in experiments [14].

In what follows, we will mostly concentrate on the one dimensional version of the model, the several variants of which addressed in the literature differ first by the choice of different bath velocity statistics $\Phi(v)$ (at rest with $\Phi(v) = \delta(v)$ [3], dichotomous [5, 7, 9] or Gaussian [6, 7, 10]) and second by the collision kernel accounting for grain interactions, that can be indexed by a continuous parameter ν (hard rods with $\nu = 1$ in most earlier works, but also $\nu = 0$ for Maxwellian rods [15] or the so-called "very hard rods" having $\nu = 2$ [10]). Moreover, in addition to mass ratio and field strength g that play a key role, collisions between the tracer and bath particles can be inelastic [16, 17]. The ensuing parameter space is therefore large, with associated rich behavior. At the Boltzmann equation level, our goal is here to provide a comprehensive view of the tracer stationary velocity distribution function, that in all generality will be denoted $f_\nu(v, g, a|\Phi)$: it depends on three parameters (ν , g and a number a that lumps mass ratio and dissipation in a single constant [8, 18]), and is furthermore a functional of bath velocity distribution $\Phi(v)$. The reason for considering an arbitrary bath distribution lies in the wide spectrum of non-equilibrium steady states that can be achieved upon driving granular gases with different energy injection mechanisms [17, 19]; Stretched exponentials $\Phi(v) \sim \exp[-|v|^\mu/\mu]$ are often encountered here.

Before rationalizing the behavior of f_ν , the model will be precisely laid in section II, where a convenient integral form will be derived from the Boltzmann equation. This provides the basis for an efficient numerical resolution algorithm, that will be described in section II C, together with the more versatile Gillespie and Direct Simulation Monte Carlo techniques. Thereby equipped with three different numerical schemes, we will put to the test in section III analytical results to be derived for the velocity tails and diffusive properties, for an arbitrary bath distribution. The cold bath setting will be put forward and solved in Section IV as a relevant model onto which more general situations can be mapped, while particular attention will be paid in Section V to the low velocity limit for bounded bath distributions. Finally, our conclusions will be presented in Section VI.

II. THE MODEL, ITS INTEGRAL REFORMULATION, AND NUMERICAL RESOLUTION

A. Statement of the problem

We consider the one dimensional motion of a single particle of mass m_t , accelerated by a constant force $g > 0$ through a bath. The bath is made up of unaccelerated particles of mass m_b , having a stationary velocity distribution function $\Phi(v)$ with characteristic velocity v_b , which is set to unity unless otherwise specified. The time dependent tracer velocity distribution is indexed by a parameter ν that specifies the type of tracer-bath interactions, and is denoted $F_\nu(v, t)$. The dependence on other parameters and on Φ is left implicit unless necessary.

The dynamics under study is governed by the linear Boltzmann (or Boltzmann-Lorentz) equation

$$\frac{\partial F_\nu(v, t)}{\partial t} + g \frac{\partial F_\nu(v, t)}{\partial v} = J[F_\nu(t), \Phi](v) \quad (1)$$

Here, J is the collision operator

$$J[\psi, \phi](v) = \int_{-\infty}^{+\infty} dv_1 dv_2 \psi(v_1) \phi(v_2) |v_1 - v_2|^\nu [\delta(v'_1 - v) - \delta(v_1 - v)] \quad (2)$$

where the post collisional velocity v'_1 of a binary encounter (v_1, v_2) can in general be written:

$$v'_1 = v_1 - (v_1 - v_2)(1 - a) = av_1 - (a - 1)v_2 \quad (3)$$

Such an expression unifies elastic and dissipative collision through the dimensionless parameter [18, 20]

$$a = \frac{m_t - \alpha m_b}{m_t + m_b} \quad (4)$$

α being the restitution coefficient which describes the inelasticity of tracer-bath collisions (elastic if $\alpha = 1$, and inelastic if $\alpha < 1$ [16]). The “memory-less” situation $a = 0$ is of particular interest due to its simplicity: each collision with a bath particle erases any memory of the tracer’s previous state, and this case has been thoroughly studied and solved in different settings [3, 6, 18]. It should also be noted that the molecular chaos assumption underlying the Boltzmann equation approach –motivated by the desire to derive analytical results– prevents the occurrence of single-file diffusion [21]. However, when possible, we will discuss systems with a higher dimensionality, for which thus the molecular chaos assumption is justified in the low density limit.

The exponent ν introduced in (2) encodes different scattering behaviors [17, 22]; The three most common models are $\nu = 1$ corresponding to Hard Rods, $\nu = 0$ for Maxwell particles and $\nu = 2$ for Very Hard Rods, the latter two approaches being often invoked to reproduce qualitatively some hard rod properties while simplifying calculations [23]. Their merits are discussed in [7] among others, but it will be shown in later sections that the value of ν may affect significant properties of the solution, some of which may exhibit crossovers depending on the value of ν .

It has been shown [10] that the velocity distribution quickly relaxes to a stationary solution $f_\nu = \lim_{t \rightarrow \infty} F_\nu$, on which we shall concentrate in the remainder: it obeys

$$g \frac{df_\nu(v)}{dv} = \int_{-\infty}^{+\infty} dv_1 dv_2 |v_1 - v_2|^\nu f_\nu(v_1) \Phi(v_2) [\delta(v'_1 - v) - \delta(v_1 - v)] \quad (5)$$

which expresses the balance between the energy received from the external field g , and the energy transferred to the bath through collisions. Eq. (5) may be rewritten as

$$g \frac{df_\nu(v)}{dv} = r_\nu(v) [H_\nu(v) - f_\nu(v)] \quad (6)$$

where

$$r_\nu(v) = \int_{-\infty}^{+\infty} du |u - v|^\nu \Phi(u) \quad (7)$$

The latter quantity is the velocity-dependent collision frequency in the gas, and $H_\nu(v)$ is given by

$$r_\nu(v) H_\nu(v) = \int dv_1 dv_2 |v_1 - v_2|^\nu f_\nu(v_1) \Phi(v_2) \delta(v'_1 - v) \quad (8)$$

which is the gain term of the collision operator, i.e. the transition rate toward velocity v through the collision process, integrated over all possible initial velocities (dummy variable v_1). Clearly, $f_\nu(v) = H_\nu(v)$ is the equilibrium solution for vanishing acceleration, when detailed balance entails that entering and leaving fluxes are equal at each point in phase space. In this field-free case ($g = 0$), two situations discussed in [18] ensure that the tracer velocity distribution is directly given by the bath distribution: first, for memory-less collisions ($a = 0$), we have

$$f_\nu(v, 0, 0|\Phi) = \Phi(v) \quad (9)$$

for arbitrary bath statistics, as can be checked from (8). Second, when the bath distribution is Gaussian $\Phi(v) = G(v) = (2\pi)^{-1/2} \exp(-v^2/2)$, one finds (see also [8])

$$f_\nu(v, 0, a|G) = G\left(\sqrt{\frac{1+a}{1-a}}v\right) \quad (10)$$

In general however, the tracer distribution differs from the bath distribution even in this unaccelerated limit, except in the tails (see section III).

We will later need to consider the distribution of pre-collisional velocities, i.e. velocities sampled right before each collision rather than uniformly over time. This distribution is usually biased toward high velocities –which have enhanced collision rate as long as $\nu > 0$ – and it is given by [24]

$$\tilde{f}_\nu(v) = \frac{r_\nu(v)}{\omega} f_\nu(v) \quad (11)$$

where the mean collision frequency is defined as

$$\omega = \int_{-\infty}^{+\infty} dv r_\nu(v) f_\nu(v) \quad (12)$$

B. Implicit form

It proves convenient to recast the stationary probability density function $f_\nu(v)$ as an implicit integral form. To this end, we write the velocity of the tracer at any given time as $v = v_1 + gt$, with v_1 the velocity that was acquired during its last encounter with a scatterer, and gt the contribution of the driving field during the timespan of ballistic flight t . We next consider the conditional probability that no collision occurs during t for a tracer starting with velocity v_1 , which involves the velocity-dependent collision frequency $r_\nu(v)$:

$$P(t|v_1) = \prod_{k=1}^{t/\Delta\tau} [1 - \Delta\tau r_\nu(v_0 + gk\Delta\tau)] \xrightarrow{\Delta\tau \rightarrow 0} \exp\left(-\int_0^t d\tau r_\nu(v_1 + g\tau)\right) \quad (13)$$

We then need the probability density for the post-collisional velocity v_1 . By definition it is proportional to the gain term of the Boltzmann operator $r_\nu(v_1) H_\nu(v_1)$, which covers all collision events from which the tracer may emerge with velocity v_1 (proper normalization of the probability densities will be enforced *a posteriori*). Finally, integrating over all possible time-spans $t \geq 0$, the stationary tracer velocity distribution can be written

$$f_\nu(v) = \int_0^{+\infty} dt \exp\left(-\int_0^t d\tau r_\nu(v - g\tau)\right) r_\nu(v - gt) H_\nu(v - gt) \quad (14)$$

It can be checked by direct calculation that the above function obeys the Boltzmann-Lorentz equation (6). The stationary solution $f_\nu(v)$ may then be interpreted as an eigenfunction of a linear integral operator \mathcal{L}_ν , associated with the eigenvalue 1:

$$f_\nu(v) = \mathcal{L}_\nu[f_\nu](v) = \int_{-\infty}^{+\infty} du f_\nu(u) K_\nu(u, v) \quad (15)$$

where the kernel $K_\nu(u, v)$ is identified from equation (14):

$$K_\nu(u, v) = (1-a)^{-\nu-1} \int_0^{+\infty} dt \exp[R_\nu(v - gt) - R_\nu(v)] |u - v + gt|^\nu \Phi\left(\frac{v - gt - au}{1-a}\right) \quad (16)$$

with

$$R_\nu(v) = \frac{1}{g} \int_0^v du r_\nu(u) \quad (17)$$

This form simplifies in the case of Maxwell particles ($\nu = 0$)

$$r_0(v) = \int du \Phi(u) = 1 \quad \text{and} \quad R_0(v) = \frac{v}{g} \quad (18)$$

The kernel then reduces to a Laplace transform

$$K_0(u, v) = \frac{1}{1-a} \int_0^{+\infty} dt e^{-t} \Phi\left(\frac{v-gt-au}{1-a}\right) \quad (19)$$

If furthermore $a = 0$, the solution itself becomes a transform of the bath distribution (as noted in [7, 10] in the special case of a Gaussian bath $\Phi(v) = G(v)$):

$$f_0(v, g, 0) = \int_0^{+\infty} dt e^{-t} \Phi(v-gt) \quad (20)$$

For general values of ν and a however, no obvious simplification can be found for the kernel $K_\nu(u, v)$. Expression (15) nevertheless allows for efficient numerical solving, as we now describe.

C. Numerical resolution of the Boltzmann-Lorentz equation

Three different techniques have been employed to obtain numerically the tracer velocity statistics. We start by summarizing their main features, before testing their compatibility.

1. Numerical integration and iterative solving

The operator \mathcal{L}_ν derives from the Boltzmann-Lorentz equation, which by construction preserves the total probability in phase space. \mathcal{L}_ν must therefore conserve the integral of any function upon which it is applied. This implies that its spectrum is reduced to two eigenspaces, associated with eigenvalues 1 (where the sought solution lies) and 0 (with non-physical eigenfunctions having vanishing integral). The application of \mathcal{L}_ν therefore acts as a projection on the eigenspace associated with the eigenvalue 1.

Numerically, discretizing \mathcal{L}_ν into a matrix L_ν and starting with a random vector, the iterative application of L_ν and subsequent normalization of the vector should converge toward the physical solution. Indeed all elements of L_ν are positive, therefore its largest eigenvalue is non-degenerate by the Perron-Frobenius theorem, associated with a positive eigenvector. This unique eigenvector coincides with the physical solution obtained through numerical simulations or determined analytically whenever possible, and this technique allows for fast computation of the solution to any precision, especially in the large velocity tails where other methods may become inefficient. At this point, we have to choose between two strategies: solving the eigenproblem numerically for the matrix L_ν , or alternatively applying the matrix iteratively until proper convergence is achieved. We have checked that both routes are equally precise and efficient, for a similar computational cost.

2. Gillespie algorithm

The second numerical technique used relies on the Gillespie algorithm, adapted to the Boltzmann-Lorentz equation by Talbot and Viot [25]. The salient features are as follows. We recall from (13) the conditional probability that no collision occurs during time t for a tracer starting with the post-collisional velocity v_1

$$\ln P(t|v_1) = R_\nu(v_1 + gt) - R_\nu(v_1) \quad (21)$$

This relationship may be inverted (either analytically or numerically by dichotomy, the latter being necessary for $\nu \neq 0, 2$ or with a non-Gaussian bath) to obtain $t(P(v_1 + gt|v_1))$, allowing us to randomly generate collisions times for the tracer, after having drawn P uniformly in $[0,1]$.

The velocity v_2 of the collision partner must then be generated with the correct weight $w(v')$. First, the flux of collisions is divided in two contributions, coming from the right and left of the tracer:

$$r_+(v_1) = \int_{-\infty}^{v_1} du |v_1 - u|^\nu \Phi(u) \quad (22)$$

$$r_-(v_1) = \int_{v_1}^{+\infty} du |v_1 - u|^\nu \Phi(u) \quad (23)$$

The probability of a collision coming from the right (resp. the left) is given by $r_+(v_1)/r(v_1)$ (resp. $r_-(v_1)/r(v_1)$). The direction of the next collision is thus determined using another random uniform number over $[0, 1]$. Depending on which side is chosen, $w(v_2) = w_\pm(v_2)$ given by

$$w_\pm(v_2) = |v_2 - v_1|^\nu \frac{\Phi(v_2)}{r_\pm(v_2)} \quad (24)$$

The velocity v' must be drawn with this weight, which is performed using an acceptance-rejection process: v_2 is generated according to the bath distribution $\Phi(v_2)$ (which significantly improves the efficiency in generating high velocities over a uniform generation with a cut-off), and then accepted with a probability

$$P_{acc}(v_2) = \frac{w_\pm(v_2) \Phi(v_0)}{\Phi(v_2) w_\pm(v_0)} \quad (25)$$

with v_0 the velocity for which $w(v_0)$ is maximal (which can be computed analytically for $\nu = 0, 1, 2$).

3. Direct Simulation Monte Carlo

We also used Direct Simulation Monte Carlo (DSMC) [26] simulations as an alternative to the Gillespie algorithm, in order to experiment easily on various baths and collision laws, without performing the (bath-dependent) analytical computations which increase the efficiency of the Gillespie method. DSMC differs from the previous approach in that instead of generating the correct collision times directly, a collision partner is drawn at each step for the tracer, with a velocity distributed according to the bath probability density function $\Phi(v_2)$. This collision event is then accepted with probability

$$P_{acc} = \frac{\Omega(v_2 - v_1)}{\Omega_{max}} \quad (26)$$

with $\Omega(v_2 - v_1) = |v_2 - v_1|^\nu$ and Ω_{max} the maximal value encountered for Ω at this point. The time counter is incremented by $1/\Omega(v_2 - v_1)$ at each step. This procedure converges in a few hundred steps per particle toward a satisfactory evaluation of the correct collision frequency, as the observed velocity distribution for the tracer (over many realizations, and/or sampled uniformly over time) is then identical to the result of the Gillespie algorithm, or analytical computations whenever they can be performed.

This method is not significantly less efficient than the Gillespie algorithm, despite its lesser specificity. It can be extended to represent collisions in higher dimensionality –as required for the discussion in section III B– by drawing a vector $\hat{\sigma}$ uniformly on the unit sphere and redefining

$$\Omega(\mathbf{v}_2 - \mathbf{v}_1) = \sigma_1^{d-1} [(\mathbf{v}_2 - \mathbf{v}_1) \cdot \hat{\sigma}]^\nu \quad (27)$$

with σ_1^{d-1} the cross-section of the tracer-bath collisions. The post collisional velocity for the tracer is then given by

$$\mathbf{v}'_1 = \mathbf{v}_1 + (1 - a) [(\mathbf{v}_2 - \mathbf{v}_1) \cdot \hat{\sigma}] \hat{\sigma} \quad (28)$$

4. Assessing the three numerical techniques

The implicit route allows for efficient computation of the stationary tracer velocity distribution with important precision. It will be below the method used whenever a specific prediction requires unusual numerical precision (good sampling of the tails). This will in particular be the case of Figs. 5 and 6. Among the three techniques, DSMC appears on the other hand to be the more versatile, since a change in the model parameter like softness exponent ν does not lead to any complication. Finally, Gillespie and DSMC algorithms are well suited to study the transient regime before the steady state is reached (see e.g. [7, 9, 10]), together with diffusive properties. Fig. 1 demonstrates the validity and consistency of all three techniques in two settings.




FIG. 1. Left: Time dependent velocity distribution $F_0(v, t, g = 1, a = 0)$, in a Gaussian bath $\Phi(v) = (2\pi)^{-1/2} \exp(-v^2/2)$ starting from a Gaussian initial condition with half variance. Gillespie simulation results (symbols) are compared to the analytic solution obtained in [7] (solid lines). Right: stationary solution $f_1(v)$, with the same parameters and initial conditions, in a Lorentzian bath $\Phi(v) = \pi^{-1}/(1 + v^2)$. The results from numerical integration (solid line) and DSMC simulation at large times (symbols) are seen to be fully consistent.

III. VELOCITY TAILS AND DIFFUSIVE BEHAVIOR

A. Asymptotics

As we shall disregard the time evolution of the system, the term "asymptotic behavior" refers here to the behavior of the stationary tracer distribution $f_\nu(v)$ for extreme values of the velocity. Rewriting the implicit equation as

$$f_\nu(v) = \int_0^{+\infty} dt \exp [R_\nu(v - gt) - R_\nu(v) + \ln H_\nu(v - gt)] r_\nu(v - gt) \quad (29)$$

with H_ν and R_ν defined in (8) and (17), we may use v as a large parameter in Laplace's method to approximate the integral over t . It is easily seen that

$$R_\nu(v) \approx \frac{|v|^{\nu+1}}{(\nu+1)g} \quad \text{when } v \rightarrow \infty \quad (30)$$

while we may choose a general stretched exponential expression for the tails of the bath distribution

$$\ln \Phi(v) \approx -|v|^\mu / \mu \quad (31)$$

where $\mu > 0$, as found in various granular systems [17, 19, 27]. Power-law tails are obtained in the limit $\mu \rightarrow 0$.

We may gain intuitive insight into the large v asymptotics by noting that competing effects are at work: such velocities may be reached either through collision with sufficiently energetic bath particles, or through acceleration without collision for a sufficiently long time. The relative importance of these effects is thus determined on the one hand by the abundance of energetic particles in the bath, characterized by the parameter μ , and on the other hand by the velocity-dependent collision frequency characterized by ν . Consequently, if ν is large enough compared to μ , the increase in collision frequency for high velocities is so steep that ballistic flight is interrupted before the tracer can be significantly accelerated. The largest velocities are therefore reached through collisions with energetic scatterers, and the Boltzmann equation is dominated by the gain term, entailing that the tracer distribution is pushed back toward the bath distribution. This may be seen as thermalization with the bath tails. If on the other hand μ is sufficiently large compared to ν , the bath tails are relatively depleted, while ballistic flight and acceleration are less impeded by the collision process. This corresponds to a prevalence of the loss term in the master equation: it is more probable for the tracer to hold a large velocity before a collision (due to acceleration) than after it.

A more refined asymptotic analysis confirms the above qualitative features. Since the argument of the exponential in (29) decreases as $t \rightarrow \infty$, it is either monotonic inside the integration interval and its maximum is located on the

boundary at $t = 0$, or it has at least one local maximum at $t = t_0$ such that

$$-g r_\nu(v - gt_0) + \frac{d \ln H_\nu}{dt}(v - gt_0) = 0 \quad (32)$$

from which we deduce that

$$t_0 = \frac{v}{g} + O(v^0) \quad (33)$$

If the maximum is located at $t = 0$, then R_ν disappears from (29) (negating the effect of the field), whereas $\ln H_\nu$ disappears at $t_0 \approx v/g$ (negating the effect of the bath), therefore the leading behavior in the positive tail is simply controlled by the value of the integrand at each of these two locations. In the negative tail, however, only one behavior prevails: when $v \rightarrow -\infty$, t_0 becomes negative and the maximum of the integrand on the domain \mathbb{R}^+ is necessarily located at $t = 0$, regardless of the parameters μ and ν . In this case we have, as in the unaccelerated problem,

$$f_\nu(v) \sim H_\nu(v) \sim \Phi(\lambda v) \quad (34)$$

where the dilation coefficient λ may be determined self-consistently using once again Laplace's method to solve $f_\nu(v) \sim H_\nu(v)$ in (8):

$$\lambda = \frac{|1 - |a|^{\mu/(\mu-1)}|^{(\mu-1)/\mu}}{1 - a} \quad \text{and} \quad \lambda = \frac{a}{1 - a} \quad \text{if} \quad \mu = 1 \quad (35)$$

For a Gaussian bath (for which $\mu = 2$) we find $\lambda^2 = (1 + a)/(1 - a)$ which equals m_t/m_b when collisions are elastic, as expected from our thermalization argument.




TABLE I. Asymptotic behaviors for the positive tail $v \rightarrow +\infty$: thermalized (left column of the table) or field-driven (right column). The negative tail $v \rightarrow -\infty$ always exhibits the thermalized behavior, see Eq. (34).

The large velocity scenario –sketched in Table I– is as follows. If $\nu + 1 > \mu$, we recover the bath behavior in $\exp(-v^\mu)$, in agreement with the intuitive argument outlined above, while if $\nu + 1 < \mu$, the positive velocity tail exhibits a $\exp(-v^{\nu+1})$ decay, hereafter referred to as the cold bath behavior. These expectations are fully corroborated by Fig. 2. If $\mu = \nu + 1$, both behaviors correspond to the same exponent, and prevalence of one process over the other depends on the value of the driving field g compared to $\lambda^{-\mu}$. For a Gaussian bath with $\mu = 2$, the borderline case corresponds to the hard-rod model, as depicted in the middle picture of Fig. 2: upon increasing the acceleration, the positive tail switches continuously from the field-independent expression $\exp[-(\lambda v)^2/2]$ when $g\lambda^2 < 1$ to the field-dependent $\exp[-v^2/2g]$ when $g\lambda^2 > 1$. The critical field, $1/\lambda^2$, reduces to unity in the memoryless case $a = 0$.

If $\nu + 1 > \mu$, the onset of the positive tail behavior described above is given by the point where

$$v^{\nu+1} \sim g\lambda^\mu v^\mu \quad (36)$$

The bath-like tail is thus expected to appear only when $v \gg v_c = (g\lambda^\mu)^{(\nu+1-\mu)^{-1}}$. Below this threshold, the acceleration usually dominates in some intermediate asymptotic range $v_b < v < v_c$ between the thermal velocity and the threshold, leading us to expect a "cold bath" behavior similar to the case $\mu \rightarrow \infty$ that will be further considered in section IV. On the left-hand side of Fig. 2, the crossover from the cold bath (field-driven) behavior to the correct tail can be seen on the curve with $v_c = g\lambda^2 = 0.5$ (crosses) while the field-driven expression remains observable for a large velocity range with a higher value of $v_c = 5$ (filled dots), for which the thermalized tail cannot be seen, but would prevail further down in the tail. The case $\mu > \nu + 1$ offers no such clear-cut crossover, as the acceleration already tends to prevail for intermediate velocities as previously discussed, and the tail behavior is therefore observed as soon as $v > v_b$. The limit $g \rightarrow 0$ is singular in this parameter range, as it suppresses the cold bath (field-driven) behavior altogether, rendering our discussion of limiting cases invalid as the only possible asymptotic behavior is the thermalized expression $f_0(v) \sim \Phi(\lambda v)$ in both tails.




FIG. 2. Asymptotic behavior in a Gaussian bath ($\mu = 2$). From left to right: $f_2(v)$, $f_1(v)$ and $f_0(v)$, computed from DSMC simulations for $a = -1/3$, $g = 1$ (crosses) and $a = 1/3$, $g = 3$ (dots). The solid line is the cold bath behavior $f_\nu(v) \propto \exp[-v|v|^\nu/(\nu+1)g]$, while the dashed line is the "equilibrium" asymptotic $f_\nu(v) \propto \Phi(\lambda v)$ which always describes the left tail ($v \rightarrow -\infty$). For $f_2(v)$ (left), the same asymptotic expression describes the right tail, although the value of the threshold $v_c = g\lambda^\mu$ may be too high for the crossover to be easily observable (as seen here with the dots which retain the cold bath behavior for the whole velocity range reported).

B. Diffusive properties

Another point of interest lies in the tracer diffusion coefficient D , which was computed in [10] for a Gaussian bath and $a = 0$, using both its Green-Kubo expression and its definition in the hydrodynamic diffusion mode of the inhomogeneous Boltzmann-Lorentz equation. These two methods were shown to give identical results for $\nu = 0$ and $\nu = 2$. The second method was also applied in [9] to the case $\nu = 1$ and $a = 0$ with a dichotomous bath, revealing the complex dependence of the diffusion coefficient on the magnitude of the field, with a minimum at finite g , while this coefficient D was shown to be monotonically increasing for $\nu = 0$ and decreasing for $\nu = 2$ in the aforementioned discussion of the Gaussian bath [10]. Here we consider an arbitrary bath distribution and parameter a , and observe once again that the effects of acceleration and thermalization disentangle when $\nu = 0$. We also demonstrate that these results are qualitatively unchanged in higher dimensional space, as reported for other observables in [8].

As shown in the appendix, letting $\nu = 0$ allows us to relate the diffusion coefficient to the variance of the tracer velocity through a simple combination

$$D = \frac{\langle v^2 \rangle - \langle v \rangle^2}{1 - a} \quad (37)$$

This variance may easily be determined recursively from that of the bath, by integration over the Boltzmann-Lorentz equation

$$\langle v^n \rangle - g n \langle v^{n-1} \rangle = \lambda^{-n} \int_{-\infty}^{+\infty} du f_0(u) \phi_n(u) \quad (38)$$

with $\phi_n(u) = \frac{1}{a-1} \int_{-\infty}^{+\infty} dv v^n \Phi\left(\frac{v-au}{a-1}\right)$ the n th moment of the displaced bath distribution. Therefore, if the bath velocity distribution is centered and has finite variance,

$$D = \frac{1}{1+a} \left(\langle v^2 \rangle_b + \frac{g^2}{(1-a)^2} \right) \quad (39)$$

where $\langle v^2 \rangle_b$ is the mean velocity squared in the bath, chosen equal to 1 in our dimensionless variables. This result is

in agreement with [10] in the limit $a \rightarrow 0$. It may be generalized to d -dimensional space (see Appendix A)

$$D_d = \frac{d}{A_d(1+a)} \left(1 + \frac{g^2 d^2}{A_d^2 (1-a)^2} \right) \quad (40)$$

where $A_d = 2\pi^{d/2}/\Gamma(d/2)$ is the surface area of a d -dimensional hypersphere. This prediction is in very good agreement with the DSMC simulation data shown in Fig. 3. In order to study the diffusive behavior of the tracer in these otherwise homogeneous simulations, its spatial coordinate is computed as the time-integral of the velocity. Between two collisions, the "space counter" in the direction of the acceleration is thus incremented by $vt + \frac{1}{2}gt^2$ with v the velocity of the tracer during this free flight and t the previously determined collision time.




FIG. 3. The diffusion coefficient D_d for $d = 1, 2, 3$ and $g = 1$ for Maxwell particles in a Gaussian bath, against the parameter a . The symbols represent DSMC simulations, while the solid lines are obtained using equation (40).

IV. THE COLD BATH

In our previous discussion of the velocity tails for the tracer in a general bath, we have observed the prevalence of a "cold bath" asymptotic behavior for certain values of the parameters, see Table I. The cold bath refers to a particular limit of the general Lorentz problem, where the scatterers are motionless and their velocity distribution is represented by the Dirac delta function: it may be understood as the limit $\mu \rightarrow \infty$ with μ defined in (31). This setting has been studied for hard rods $\nu = 1$ with $a = 0$ [3] and in some limiting cases with $a \leq 0$ [8]. Inserting $\Phi(v) = \delta(v)$ in (7) gives $r_\nu(v) = |v|^\nu$ and the Boltzmann-Lorentz equation takes the form of a functional-differential equation

$$gf'_\nu(v, g, a) = |v|^\nu \left(\frac{1}{|a|^{\nu+1}} f_\nu\left(\frac{v}{a}, g, a\right) - f_\nu(v, g, a) \right) \quad (41)$$

which is reminiscent of relations found in other stochastic processes such as random walks and growth models on trees [28]. In the field-free case, this equation reverts to a class of random collision problems systematically studied by ben-Avraham et al. [29].

This system plays a prototypical role in the study of the general Lorentz problem for two reasons. First, for Maxwell particles ($\nu = 0$), a theorem proved by Wannier [30] for the three-dimensional case and Eder and Posch [7] in one dimension gives the complete (time-dependent) solution for any bath distribution Φ as a convolution over the cold bath solution

$$F_0(v, t, g, a|\Phi) = \int_{-\infty}^{+\infty} du \Phi(u) F_0(v-u, t, g, a|\delta) \quad (42)$$

This relation holds as well for the stationary state $f_0(v)$ which is under scrutiny here.

On the other hand, given its expression for $\nu = 0$, we may derive the cold bath solution for any value of the collision kernel exponent. This may be seen if we define

$$f_\nu(v) = h_0 \left(\frac{v|v|^\nu}{\nu+1} \right) \quad (43)$$

which verifies

$$gh'_0(v) = \frac{1}{|a|^{\nu+1}} h_0\left(\frac{v}{a|a|^\nu}\right) - h_0(v) \quad (44)$$

With the simple rescaling $a|a|^\nu \rightarrow a$ this equation is made to coincide with (41), which is associated to Maxwell particles.

Therefore, the cold bath solution allows for direct computation of solutions that fall either on the vertical axis $\nu = 0$ or on the horizontal line $\mu = \infty$ on the graph presented beside Table I. However Wannier's convolution theorem does not hold for arbitrary values of ν , which prevents us from extending the cold bath solution to settings where both ν and μ are chosen different from the prototypical case ($\nu = 0, \mu = \infty$). Yet, this solution may still preserve significant features of the general solution, such as the positive tail behavior in the conditions discussed in section III.

A. Explicit solution

The cold bath equation (41) may be rewritten as the implicit equation

$$f_\nu(v) = \frac{1}{g} \int_{-\infty}^{+\infty} du f_\nu(u) |u|^\nu \exp\left[-\frac{1}{g(\nu+1)}(v|v|^\nu - au|au|^\nu)\right] \Theta(v - au) \quad (45)$$

where Θ denotes the Heaviside function. For finite a , we recall the characteristic "cold bath" positive tail behavior presented in Table I: when $v \rightarrow +\infty$,

$$f_\nu(v) \propto \exp\left[-\frac{v^{\nu+1}}{g(\nu+1)}\right] \quad (46)$$

The negative tail behavior however differs from the situations discussed in section III, where it was determined by the bath distribution. As we shall see, the negative tail exists in the cold bath only if $a < 0$. In this case, the term $f_\nu(v/a)$ which appears in (41) is located in the positive tail, allowing us to self-consistently neglect the loss term and expect

$$f_\nu(v \rightarrow -\infty, g, a < 0) \propto \exp\left[-\frac{(v/a)^{\nu+1}}{g(\nu+1)}\right] \quad (47)$$

These asymptotic expressions for $|v| \rightarrow \infty$ with $a < 0$ are corroborated by numerical results in Fig. 4.




FIG. 4. Asymptotic behavior for a Maxwellian tracer in a cold bath and a Gaussian bath ($a = -1/3, g = 3$). The symbols are Gillespie simulation results, the solid line is the exact cold bath solution (59), and the dashed lines represent the asymptotic slopes $\exp(-v/g)$ (right, see Eq. (46)) and $\exp(-v/ag)$ (left, see Eq. (47)). Even at finite temperature, a cold bath-like behavior dominates for a large velocity range in both tails (although the negative tail eventually becomes Gaussian, as clearly seen with different values of g and a in Fig. 2).

It should be emphasized here that the negative tail $v \rightarrow -\infty$ does not exist whenever $a \geq 0$: in that case, equation (3) entails that the tracer velocity is reduced but never reverted by collisions with bath particles. Once it has become

positive due to acceleration, it may not become negative again, which ensures that the stationary solution has support on \mathbb{R}^+ only. This feature –which will be further developed in section V– indeed is apparent in the case $a = 0$, which is straightforwardly derived from (45) (note the choice of index $\nu - 1$ for simpler expressions):

$$f_{\nu-1}(v, g, 0) = \Theta(v) \frac{\nu}{(\nu g)^{\nu-1} \Gamma(\nu-1)} \exp\left[-\frac{v^\nu}{\nu g}\right] \quad (48)$$

where it appears that the solution has support on $v > 0$. Another interesting property of this solution is the mean velocity

$$\langle v \rangle_{\nu-1} = \frac{1}{2\sqrt{\pi}} \Gamma\left(\frac{2+\nu}{2\nu}\right) (4\nu g)^{1/\nu} \quad (49)$$

where Γ is the Euler gamma function. Thus $\langle v \rangle_\nu \propto g^{1/(\nu+1)}$ in agreement with the dimensional argument presented in [3]. It should be noted that, unless $\nu = 0$, this implies a breakdown of linear response: for $\nu = 1$, $\langle v \rangle \propto g^{1/2}$ as discussed in [8].

We now derive the full solution for $a \neq 0$, considering Maxwellian particles (without loss of generality, as argued above) which obey

$$g f'_0(v) = \frac{1}{|a|} f_0\left(\frac{v}{a}\right) - f_0(v) \quad (50)$$

The moment hierarchy is obtained by a straightforward integration, and its general term is found

$$g n \langle v^{n-1} \rangle = \langle v^n \rangle (1 - a^n) \quad (51)$$

$$\langle v^n \rangle = n! g^n b_n \quad (52)$$

where $b_n = 1/(a, a)_n$ using the q -Pochhammer notation [31]

$$(a, q)_n = \begin{cases} 1 & n = 0 \\ \prod_{m=0}^{n-1} (1 - a q^m) & n \geq 1 \end{cases} \quad (53)$$

This symbol appears in combinatorics as a generalization of the Pochhammer symbol or rising factorial:

$$\lim_{q \rightarrow 1^-} \frac{(q^x, q)_n}{(1 - q)^n} = x(x+1)\dots(x+n-1) = \frac{\Gamma(x+n)}{\Gamma(x)} \quad (54)$$

The expression $(a, a)_n$ converges in the limit $n \rightarrow \infty$ to a standard function of a known as the Euler function (not to be confused with the better known Euler totient function in number theory). The general expression of the solution may be found by taking the Fourier transform of the function

$$\hat{f}_0(s) = \int_{-\infty}^{+\infty} f_0(v) e^{isv} dv \quad (55)$$

which in turn verifies

$$igs \hat{f}_0(s) = \hat{f}_0(sa) - \hat{f}_0(s) \quad (56)$$

$$\hat{f}_0(s) = \frac{\hat{f}_0(sa)}{1 + igs} \quad (57)$$

Then by recurrence and using $\hat{f}_0(0) = 1$ due to the normalization of $f_0(v)$

$$\hat{f}_0(s) = \lim_{n \rightarrow \infty} \frac{\hat{f}_0(sa^{n+1})}{\prod_{k=0}^n (1 + igs a^k)} = \prod_{k=0}^{\infty} (1 + igs a^k)^{-1} \quad (58)$$

Taking the inverse transform (with some care as the position of the poles and the integration contour depend on the sign of both a and v)

$$f_0(v) = \sum_{k=0}^{\infty} \frac{e^{-v/ga^k}}{ga^k} \Theta(v/ga^k) \prod_{j \geq 0, j \neq k} (1 - a^{j-k})^{-1} \quad (59)$$

From this expression we may verify $f_0(0) = 0$ if $a > 0$, and identify the asymptotic behavior for $|v| \rightarrow \infty$ as the first non-zero term in the sum depending on the sign of v and a , see Fig. 4. Furthermore, we may rewrite the product over j using the b_n notation defined in (53), while the exponential term is recognized as the solution for $a = 0$ in (48) where the acceleration has undergone the rescaling $g \rightarrow ga^k$. We obtain the concise formula

$$f_0(v, g, a) = b_{\infty} \sum_{k=0}^{\infty} (-1)^k a^{k(k+1)/2} b_k f_0(v, ga^k, 0) \quad (60)$$

This structure is interesting in several respects: first, the sum converges very rapidly for the needs of numerical computation, and is found in excellent agreement with our simulation results. Second, due to the simple form of $f_0(v, g, 0)$, using Wannier's theorem to derive the full solution for Maxwell particles in any bath will only involve taking Laplace transforms of the bath distribution. An application to the Gaussian bath is given below. Finally, even though this structure will not be preserved for systems with arbitrary bath distribution and $\nu \neq 0$, it suggests that some significant traits of the general solution may be contained in the limiting case $a = 0$, which is more readily solved in any model, see e.g. [6, 10]. However, this analytical expression sheds little light on the behavior of the cold bath solution for positive a in the limit $v \rightarrow 0$, which will warrant a separate investigation in section V.

B. Application to the Gaussian bath

We now revisit some properties of one of the most extensively studied settings, the Maxwellian tracer in a Gaussian bath ($\nu = 0$, $\mu = 2$) [6, 7, 10]. This will serve to demonstrate the application of results from the cold bath limit to arbitrary bath distributions, which we recall is exact for $\nu = 0$.

From Wannier's convolution theorem we deduce that the solution $f_0(v, g, 0)$ in this setting is simply given by the Laplace transform of the Gaussian $G(v)$, as we previously saw in (20) (see also [10])

$$f_0(v, g, 0|G) = \frac{1}{2g} \exp\left[\frac{1}{2g^2} - \frac{v}{g}\right] \left(1 + \operatorname{erf}\left[\frac{v - g^{-1}}{\sqrt{2}}\right]\right) \quad (61)$$

Inserting this expression in the series (60) allows for accurate numerical computation as well as asymptotic analysis for $a \neq 0$.

We thus recover a result that was obtained in Ref. [7], where use was made of a method specific to this choice of bath and collision kernel: using Mehler's formula [32], it is possible to compute the analytical solution of the Boltzmann-Lorentz equation as an expansion in terms of the Hermite Polynomials $H_n(v)$

$$f_0(v|G) = G(\lambda v) \sum_{n=0}^{\infty} \left(\frac{\lambda g}{\sqrt{2}}\right)^n b_n H_n\left(\frac{\lambda v}{\sqrt{2}}\right) \quad (62)$$

However, this expansion diverges for any value of the parameters, even at vanishing velocity as $H_n(0) = (-1)^{n/2} n!/(n/2)!$ for even n , while $H_n(0) = 0$ for odd n . Consequently, the expansion must be put under a different form to allow for numerical computation. It is suggested in [7] that, owing to a theorem by Euler

$$b_n = b_{\infty} \sum_{k=0}^{\infty} (-1)^k a^{nk} a^{k(k+1)/2} b_k \quad (63)$$

from which, interchanging the sums, one finds as expected

$$f_0(v, g, a) = b_{\infty} \sum_{k=0}^{\infty} (-1)^k a^{k(k+1)/2} b_k f_0(v, ga^k, 0) \quad (64)$$

Finally, we may be interested in seeing how exactly the hierarchy of moments with a Gaussian bath relates to its cold bath equivalent. We compute the moment-generatrix, using the connection between the Hermite polynomials and the derivatives of the Gaussian

$$M(t) = \int_{-\infty}^{+\infty} dv e^{tv} f_0(v) = \exp\left[\frac{t^2}{2\lambda^2}\right] \sum_{j=0}^k \binom{k}{j} \zeta^{(k-j)}(t) \left(i\sqrt{2}\lambda\right)^{-j} H_j\left(\frac{it}{\sqrt{2}\lambda}\right) \quad (65)$$

with

$$\zeta(t) = \sum_{n=0}^{\infty} t^n g^n b_n \quad (66)$$

from which we derive a general expression for the k -th moment of $f_0(v)$:

$$\begin{aligned} \langle v^k \rangle &= \frac{d^k M}{dt^k}(t=0) \\ &= \sum_{j=0}^{\lfloor k/2 \rfloor} \frac{k!}{j!} (2\lambda^2)^{-j} g^{k-2j} b_{k-2j} \end{aligned} \quad (67)$$

The term $j=0$ (which is the leading order for large g) is recognized as the corresponding moment in the cold bath (52), while the following terms appear to be specific to the Gaussian setting.

V. NON-NEGATIVE STATIONARY DISTRIBUTION

The solution for $a > 0$ exhibits some peculiar properties in the cold bath configuration: the tracer velocity can be reduced but never reverted by a collision. Due to the positive acceleration, if the tracer reaches a positive velocity at a given time, it will subsequently never be able to revert its direction of motion. As mentioned in section IV A, this means that the stationary distribution has support on the positive semi-axis only. This gives a new limit to consider, that of small velocities $v \rightarrow 0$, which has no equivalent in any of the settings previously considered, and may appear only when the bath distribution is bounded from below.

A. Heuristic argument

We shall now show that the asymptotic behavior of $f_0(v)$ for vanishing velocities is of log-normal form. We start with a heuristic argument. In order to reach a very low velocity v , numerous collisions must happen in a very short time span, before the acceleration can restore the velocity to its typical scale. Each collision multiplies the velocity of the tracer by a , therefore a sequence of repeated collisions may be viewed as a multiplicative random walk, or an additive random walk on $\ln v$: $\ln v_{i+1} \approx \ln v_i + \ln a$ where v_i is the velocity of the tracer following collision i . After a random but large number of steps (collisions occurring almost instantaneously), we expect $\ln v$ to be normally distributed.

This intuition may be put on more solid grounds. In any sequence that allows the tracer to reach a very low velocity, collision $i+1$ must take place in a time interval Δt_i short enough that $g\Delta t_i \ll v_{i-1}$, lest the tracer be reaccelerated to its former velocity. This interval, the upper bound for the time span between two successive collisions, becomes smaller as lower velocities are involved, therefore we may choose a bound of the form $g\Delta t_i = \epsilon v_i$ with a constant factor $\epsilon \ll 1$. At least one collision must occur during each interval Δt_i , which has probability $1 - \exp(-\omega\Delta t_i) \approx \omega\Delta t_i$. Starting from a characteristic velocity \bar{v} , we need n such collisions to reach the velocity $v_n = a^n \bar{v}$, and we may associate to this sequence the following probability density

$$f_0(v_n) \propto \prod_{i=0}^n \omega\Delta t_i \approx \left(\frac{\epsilon\omega}{g}\right)^n a^{n(n+1)/2} \quad (68)$$

Writing $\chi = 1/\ln a$, we have

$$f_0(v) \sim \exp\left\{\frac{\chi}{2} \ln^2(v)\right\} \quad (69)$$

We cannot expect to obtain more than the leading order in $\ln^2(v)$ from such simple considerations, but Fig. 5 shows that the above expansion indeed provides the dominant behavior of the stationary velocity distribution as $v \rightarrow 0$, although increasingly difficult to evidence as a increases.




FIG. 5. Plot of $\ln f_0(v)/\ln^2(v)$ following from (59), as a function of $\ln^2(v)$ (increasing as $v \rightarrow 0$), for different values of a and $g = 1$. To each set is associated a horizontal line marking the expected limit $\chi/2 = (2 \ln a)^{-1}$, see Eq. (69), from which we may see the approach toward the limit becoming extremely slow as a increases.

B. Log-normal characteristic approximation

We now wish to improve upon the heuristic asymptotics (69), through a more formal derivation. It proves convenient to redefine $\hat{f}_0(s)$ as the Laplace transform

$$\hat{f}_0(s) = \int_0^{+\infty} f_0(v) e^{-sv} dv \quad (70)$$

which, for imaginary arguments, coincides with the previous definition of $\hat{f}_0(s)$ given in section IV A, and verifies again

$$\hat{f}_0(s) = \frac{\hat{f}_0(sa)}{1 + gs} \quad (71)$$

Iterating this relation, we obtain

$$\hat{f}_0\left(\frac{1}{g}\right) = \hat{f}_0\left(\frac{1}{ga^{n+1}}\right) a^{-(n+1)(n+2)/2} \prod_{k=1}^{n+1} (1 + a^k) \quad (72)$$

The product over k may be rewritten as $(-a, a)_{n+1}$ using the q -Pochhammer symbol defined in (53), and it is bounded from above by the finite asymptotic value $(-a, a)_\infty$ given as a function of a in [33]. We may thus take the limit $n \rightarrow \infty$, for which $1/ga^{n+1} \rightarrow \infty$. However $\hat{f}_0(1/g)$ is finite whereas

$$\lim_{s \rightarrow +\infty} s \hat{f}_0(s) = f_0(0) = 0 \quad (73)$$

For $\hat{f}_0(1/g)$ to be finite as $n \rightarrow \infty$, we must have

$$\hat{f}_0\left(\frac{1}{ga^{n+1}}\right) \propto a^{(n+1)(n+2)/2} \quad (74)$$

Therefore, taking $s = a^{-(n+3/2)}$, we find

$$\lim_{s \rightarrow \infty} s \hat{f}_0(s\sqrt{a}/g) \propto \exp\left[\frac{\chi \ln^2(s)}{2}\right] \quad \text{with } \chi = (\ln a)^{-1} < 0 \quad (75)$$

We emphasize here that in deriving the above asymptotic expansion, everything amounts to neglecting the loss term in the Boltzmann equation (50). Indeed, if we expect $f_0(v)$ to decrease steeply for vanishing velocities, it is much less probable to hold velocity $v \ll 1$ than v/a , allowing us to approximate (50) by

$$gf'_0(v) \approx \frac{1}{a} f_0\left(\frac{v}{a}\right) \quad (76)$$

hence for the Laplace transform

$$gs\hat{f}_0(s) \approx \hat{f}_0(as) \quad (77)$$

which may be iterated as in (72), leaving out the factor $(-a, a)_{n+1}$ but giving the same asymptotic condition (74). This equation is verified exactly by the log-normal law Ψ :

$$gs\Psi'(s) = \Psi(as) \quad (78)$$

$$\Psi(s) = \frac{1}{s\sqrt{2\pi\sigma}} \exp\left[-\frac{(\ln s - m)^2}{2\sigma^2}\right] \quad (79)$$

where m and σ are easily determined from the previously known parameters

$$\sigma^2 = -\ln a \quad (80)$$

$$m = \frac{1}{2} \ln a - \ln g \quad (81)$$

As discussed in Leipnik [34] and Lopez [35], equation (78) and its counterpart in velocity space (76) are verified exactly by multiple families of functions. We note in passing that the existence of these different families of solutions is linked with well-known indeterminacy problems surrounding the log-normal: most notable is the non-uniqueness of its moments, shared by several families of distributions, a problem studied since Stieltjes' memoir "Recherches sur les fractions continues" [36] and his discovery that a probability distribution need not be uniquely determined by its moments.

Assuming that $\hat{f}_0(s)$ is indeed of log-normal form in the limit $s \rightarrow \infty$, we may derive the corresponding asymptotic expression of $f_0(v)$ for vanishing velocities, which later receives numerical confirmation. In the spirit of [34, 37], this expression is found from an integral expression verifying (76)

$$\varphi_S(v) = \int_{k-i\infty}^{k+i\infty} \exp\left[\frac{\sigma^2 z^2}{2} - z(\ln v + m)\right] S(z)\Gamma(z) dz \quad (82)$$

where $S(z)$ is any anti-periodic function of anti-period 1: $S(z-1) = -S(z)$. Leipnik shows in [34] that the choice $S(z) = (2\pi)^{-1} \sin(\pi z)$ – that we also adopt below – equates $\varphi_S(t)$ with the characteristic function (i.e. Fourier transform) of the log-normal law. Applying Laplace's method to the integral, we expect for large $|\ln v|$

$$\varphi_S(v) \approx \exp\left[\frac{\sigma^2 z_0^2}{2} - z_0(\ln v + m)\right] S(z_0)\Gamma(z_0) \quad (83)$$

with z_0 defined as the point where the logarithmic derivative of the integrand vanishes

$$0 = \sigma^2 z_0 - (\ln v + m) + \frac{S'(z_0)}{S(z_0)} + \frac{\Gamma'(z_0)}{\Gamma(z_0)} \quad (84)$$

we may therefore take $z_0 \approx (\ln v + m)/\sigma^2$ as the other terms grow at most logarithmically. Letting $y = \ln v + m$ suggests the asymptotic behavior

$$\begin{aligned} f_0(v = e^{y-m}) &\propto \exp\left[-\frac{y^2}{2\sigma^2}\right] S(y/\sigma^2)\Gamma(y/\sigma^2) \\ f_0(v) &\propto \exp\left[\chi y \left(\frac{y}{2} + 1 - \ln(\chi y)\right)\right] \quad \text{for } y \equiv \ln(v\sqrt{a}/g) \rightarrow -\infty \end{aligned} \quad (85)$$

where we used $\chi \ln v \gg 1$ and Leipnik's choice $S^{-1}(z) = 2\Gamma(z)\Gamma(1-z)$ [34]. We thus recover the leading order of the heuristic expression (69), with additionally a subleading contribution $(\chi \ln v)^{-\chi \ln v}$ in the probability distribution, stemming from the gamma function. Although subleading, this correction is nevertheless important: the complete structure of expression (85) is required so that, upon insertion in the Boltzmann equation without loss term, both sides may be matched for vanishing v . Indeed, it verifies relation (76) to leading order in $v/\ln v$ whereas our first log-normal approximation (69) does not.

This new asymptotic expression is clearly seen in Fig. 6 (left) to approximate the solution $f_0(v)$ much more closely and on a wider range of velocities than (69). Furthermore the right-hand graph demonstrates the same behavior for a different non-negative bath distribution: a half-Gaussian, which strictly vanishes for $v < 0$. This tail behavior appears for $a > 0$ as long as there is a lower bound to the bath particle velocity. In addition, any distribution $\Phi(v) \neq \delta(v)$ but bounded from below with support in $[0, \infty[$ leads to a decrease of tracer velocity probabilities (see the half-Gaussian results in Fig. 6, well below their cold bath counterpart), shifting the tail by a multiplicative constant without affecting its functional form. However, the range of validity of this asymptotic expression is relegated further in the tail when a increases.




FIG. 6. Left: Validity of the asymptotic expressions. The exact cold bath solution $f_0(v)$ [solid lines, corresponding to (59)] against the predictions (69) (dashed lines) and (85) (dots). Here, $a = 0.01, 0.1$ and 0.5 from left to right, and $g = 1$. As a increases, the predicted asymptotic behavior appears further in the tail, and the expansion (85) becomes significantly more precise than our first heuristic guess (69). Right: Close-up on the data for $a = 0.01$ (solid line) and comparison with the solution corresponding to a half-Gaussian bath $\Phi(v) = \Theta(v)(2\pi)^{-1/2} \exp(-v^2/2)$ (dashed line), other parameters being identical. The same asymptotic form (85) (dots) holds in both cases. Similar behavior is expected for any bath distribution with a lower velocity bound, provided $g > 0$ and $a > 0$.

VI. CONCLUSIONS

Within a generalized Boltzmann-Lorentz framework, we have studied the stationary state of a tracer particle immersed in a bath with arbitrary velocity statistics $\Phi(v)$. The tracer (say a charged particle) is accelerated by an external field, that does not act on the bath (say made up of neutral particles). In addition to Φ , the model is specified by the field intensity g , a parameter a accounting for material properties that combines mass ratio (tracer over bath particle) together with collisional dissipation, and an index ν quantifying the softness of particle interactions. Maxwellian, hard, and very hard rods are thereby embedded in a unifying approach, that lends itself to analytical progress and numerical investigation. Three independent simulation techniques were used to solve the Boltzmann-Lorentz equation. We have performed an asymptotic analysis of the high-energy tails. We have evidenced a winner-takes-all competition between the two processes that lead to a non equilibrium stationary state: on the one hand the acceleration during the variable timespan of ballistic progress, which feeds energy into the system, and on the other hand, the collision with scatterers that lead to dissipation. Our analysis has revealed that, depending on the parameter and velocity range that we consider, either of these effects dominates and completely determines the local shape of the distribution. Therefore, significant portions of this distribution are almost independent either from the bath or from the acceleration and collision kernel. The latter effect may provide a way to probe non-gaussianities in the bath using a more visible tracer and charting out its large velocities behavior; Such a behavior requires that the scattering exponent ν is large enough once Φ has been chosen, or conversely that the bath distribution is not too cold –i.e. not too peaked around the origin– once the scattering exponent is fixed. Whereas most of the analysis was performed for a one dimensional system, we have, for diffusive properties, also considered driven discs or spheres in higher dimension.

Motivated by the existence of bath-independent properties, particular emphasis was put on an already introduced setting, but previously unsolved in its general formulation: the “cold gas” with vanishing temperature [i.e. $\Phi(v) = \delta(v)$]. Its interest as an approximation for more realistic models was addressed. We have in particular shown that even when the cold bath asymptotics does not prevail, an intermediate asymptotic cold bath regime may exist, when the tracer velocity is significantly larger than the thermal velocity v_b but lower than the model-dependent threshold v_c . This setting also allowed us to exhibit an interesting asymptotic behavior at low velocities for a tracer in a bath with a non-negative velocity distribution –e.g. a heavy intruder falling in a static bath or along a stationary stream of particles.

ACKNOWLEDGMENTS

We would like to thank A. Alastuey, A. Burdeau, S. Majumdar, J. Talbot, and P. Viot for useful discussions.

-
- [1] H. Lorentz, in *KNAW, Proceedings*, Vol. 7 (1905) pp. 1904–1905.
[2] J. Piasecki and E. Wajnryb, *J. Stat. Phys.* **21**, 549 (1979).
[3] J. Piasecki, *J. Stat. Phys.* **30**, 185 (1983).
[4] P. Résibois and M. Mareschal, *Physica A* **94**, 211 (1978).
[5] J. Piasecki, *Phy. Lett. A* **114**, 245 (1986).
[6] A. Gervois and J. Piasecki, *J. Stat. Phys.* **42**, 1091 (1986).
[7] O. Eder and M. Posch, *J. Stat. Phys.* **52**, 1031 (1988).
[8] P. Martin and J. Piasecki, *Europhys. Lett.* **46**, 613 (1999).
[9] J. Piasecki and R. Soto, *Physica A* **369**, 379 (2006).
[10] A. Alastuey and J. Piasecki, *J. Stat. Phys.* **139**, 991 (2010).
[11] G. Gradenigo, A. Puglisi, A. Sarracino, and U. M. B. Marconi, *Phys. Rev. E* **85**, 031112 (2012).
[12] A. Puglisi, A. Sarracino, G. Gradenigo, and D. Villamaina, *Gran. Matt.* (2012), 10.1007/s10035-012-0312-9.
[13] A. Fiege, M. Grob, and A. Zippelius, *Gran. Matt.* (2012), 10.1007/s10035-011-0309-9.
[14] R. Candelier and O. Dauchot, *Phys. Rev. E* **81**, 011304 (2010).
[15] J. Maxwell, *Phil. Trans. R. Soc* **157**, 49 (1867).
[16] N. Brilliantov and T. Pöschel, *Kinetic Theory of Granular Gases* (Oxford Graduate Texts, 2004).
[17] M. Ernst, E. Trizac, and A. Barrat, *J. Stat. Phys.* **124**, 549 (2006).
[18] A. Santos and J. Dufty, *Phys. Rev. Lett.* **97**, 58001 (2006).
[19] J. M. Montanero and A. Santos, *Gran. Matt.* **2**, 53 (2000).
[20] J. Piasecki, J. Talbot, and P. Viot, *Physica A* **373**, 313 (2007).
[21] K. Hahn, J. Kärger, and V. Kukla, *Phys. Rev. Lett.* **76**, 2762 (1996).
[22] M. Ernst, *Phys. Rep.* **78**, 1 (1981).
[23] P. L. Krapivsky and C. Sire, *Phys. Rev. Lett.* **86**, 2494 (2001).
[24] P. Visco, F. van Wijland, and E. Trizac, *Phys. Rev. E* **77**, 041117 (2008).
[25] J. Talbot and P. Viot, *J. Phys. A: Math. Gen.* **39**, 10947 (2006).
[26] G. Bird, *Computers & Mathematics with Applications* **35**, 1 (1998).
[27] M. Ernst and R. Brito, *Phys. Rev. E* **65**, 040301 (2002).
[28] D. Dean and S. Majumdar, *J. Stat. Phys* **124**, 1351 (2006).
[29] D. Ben-Avraham, E. Ben-Naim, K. Lindenberg, and A. Rosas, *Phys. Rev. E* **68**, 050103 (2003).
[30] G. Wannier, *Phys. Rev.* **83**, 281 (1951).
[31] G. Gasper and M. Rahman, “Basic hypergeometric series,” in *Encyclopedia of Mathematics and its Applications*, Vol. 35 (Cambridge University Press, Cambridge, 1990).
[32] H. Bateman, A. Erdélyi, and B. M. Project, *Higher transcendental functions* (Robert E. Krieger, 1981).
[33] B. Gordon and R. Mcintosh, *Journal of the London Mathematical Society* **62**, 321 (2000).
[34] R. Leipnik, *The Journal of the Australian Mathematical Society. Series B. Applied Mathematics* **32**, 327 (1991).
[35] M. López-García, *Theory of Probability and its Applications* **55**, 303 (2011).
[36] T. Stieltjes and G. Dijk, *Collected Papers/Oeuvres Completes* (Springer Verlag, 1993).
[37] N. De Bruijn, *Indagationes Math* **15**, 449 (1953).
[38] P. Résibois and M. De Leener, *Classical kinetic theory of fluids* (Wiley New York, 1977).

APPENDIX: COMPUTATION OF THE DIFFUSION COEFFICIENT

Let us define the velocity autocorrelation function $\Gamma(t)$ as

$$\Gamma(t) = \langle (v(t) - \langle v \rangle)(v(0) - \langle v \rangle) \rangle \quad (86)$$

where $\langle . \rangle$ denotes the mean over the stationary distribution $f_\nu(v)$ and $v(t)$ is the velocity of a given realization of the tracer at time t . A standard relation [3, 38] connects $\Gamma(t)$ to the diffusion coefficient through

$$D = \int_0^\infty dt \Gamma(t) \quad (87)$$

In a derivation similar to the one used in [38], this function $\Gamma(t)$ is expressed as

$$\Gamma(t) = \int dv_0 f_\nu(v_0) \int dv f_\nu(v, t|v_0) (v - \langle v \rangle)(v_0 - \langle v \rangle) \quad (88)$$

where $f_\nu(v, t|v_0)$ is the conditional velocity distribution of the tracer at time t knowing it had velocity v_0 at time 0 (it is the time-dependent distribution $f_\nu(v, t)$ with the initial condition $f_\nu(v, 0) = \delta(v - v_0)$). We may thus define the auxiliary function

$$N(v, t) = \int dv_0 f_\nu(v_0) f_\nu(v, t|v_0) (v_0 - \langle v \rangle) \quad (89)$$

which fulfills the initial condition

$$N(v, 0) = (v - \langle v \rangle) f_\nu(v) \quad (90)$$

and

$$\Gamma(t) = \int dv (v - \langle v \rangle) N(v, t) \quad (91)$$

Due to the linearity of the Boltzmann-Lorentz equation, $N(v, t)$ follows the same equation as $F(v, t)$ as can be seen from Eq. (89). Therefore, the determination of $\Gamma(t)$ using (91) amounts to computing the integral and first moment of the solution of the Boltzmann-Lorentz equation with the non-physical initial condition (90). As this equation conserves the normalization,

$$\int dv N(v, t) = \int dv N(v, 0) = 0 \quad (92)$$

thus

$$\Gamma(t) = \int dv v N(v, t) \quad (93)$$

Finally, if $\nu = 0$, this time-dependent first moment can be computed directly from the Boltzmann equation

$$\frac{\partial \Gamma}{\partial t} = -g \int dv v \frac{\partial N}{\partial v} + \int dv_1 dv_2 N(v_1) \Phi(v_2) \int dv v [\delta(v'_1 - v) - \delta(v_1 - v)] \quad (94)$$

The first term on the right-hand side vanishes, while $v'_1 - v_1 = (a - 1)(v_1 - v_2)$ and therefore

$$\frac{\partial \Gamma}{\partial t} = -(1 - a)\Gamma(t) \quad (95)$$

$$\Gamma(t) = \Gamma(0) e^{-(1-a)t} \quad (96)$$

from which the diffusion coefficient follows

$$D = \frac{\Gamma(0)}{1 - a} = \frac{\langle v^2 \rangle - \langle v \rangle^2}{1 - a} \quad (97)$$

We have thus related D to the variance of the tracer stationary velocity distribution in the case of Maxwell particles with an arbitrary bath.

Finally, the approach can be generalized to higher dimensions, where the collision law reads

$$\mathbf{v}'_1 = \mathbf{v}_1 + (1 - a) [(\mathbf{v}_2 - \mathbf{v}_1) \cdot \hat{\sigma}] \hat{\sigma} \quad (98)$$

so that equation (94) becomes

$$\frac{\partial \Gamma}{\partial t} = - \int d\mathbf{v} \mathbf{v} (\mathbf{g} \cdot \nabla_{\mathbf{v}}) \mathbf{N}(\mathbf{v}, t) + \int d\mathbf{v}_1 d\mathbf{v}_2 d\hat{\sigma} \mathbf{N}(\mathbf{v}_1) \Phi(\mathbf{v}_2) \int d\mathbf{v} \mathbf{v} [\delta(\mathbf{v}'_1 - \mathbf{v}) - \delta(\mathbf{v}_1 - \mathbf{v})] \quad (99)$$

Then, if we define \mathbf{e}_{12} the unit vector along $(\mathbf{v}_2 - \mathbf{v}_1)$, θ the angle between $\hat{\sigma}$ and \mathbf{e}_{12} , and $\hat{\Omega}$ the solid angle

$$\int d\hat{\sigma} [(\mathbf{v}_2 - \mathbf{v}_1) \cdot \hat{\sigma}] \hat{\sigma} = \int d\hat{\Omega} |\mathbf{v}_2 - \mathbf{v}_1| \cos^2(\theta) \mathbf{e}_{12} = C(d) (\mathbf{v}_2 - \mathbf{v}_1) \quad (100)$$

with $C(d)$ a dimension-dependent coefficient which is related to A_d the area of the d -dimensional unit hypersphere:

$$C(d) = \int d\hat{\Omega}[1 - \sin^2(\theta)] = A_d \left[1 - \frac{\int_0^\pi d\theta \sin^d \theta}{\int_0^\pi d\theta \sin^{d-2} \theta} \right] = \frac{A_d}{d} \quad (101)$$

We may then compute the first and second moments: letting $\mathbf{g} = g\mathbf{e}_x$

$$\langle \mathbf{v} \rangle = \langle v_x \rangle \mathbf{e}_x = \frac{\mathbf{g}d}{A_d(1-a)} \quad (102)$$

$$\langle \mathbf{v}^2 \rangle = \frac{1-a}{1+a} \langle v^2 \rangle_b + \frac{2gd}{1-a^2} \frac{\langle v_x \rangle}{A_d} \quad (103)$$

hence

$$\Gamma(0) = \langle \mathbf{v}^2 \rangle - \langle \mathbf{v} \rangle^2 = \frac{1-a}{1+a} \langle v^2 \rangle_b + \frac{g^2 d^2}{A_d^2(1-a^2)} \quad (104)$$

which gives the following relation for the diffusion coefficient in d dimensions in a bath with unit temperature:

$$D_d = \frac{d}{A_d(1+a)} \left(1 + \frac{g^2 d^2}{A_d^2(1-a)^2} \right) \quad (105)$$